


Electrochemistry

"Redox" or oxidation-reduction reactions involve a change in the oxidation state of the chemical species involved in a chemical reaction. (Review handout on oxidation numbers.)


Example:



The two **half-cell** reactions combine to yield the overall reaction. The number of electrons gained in the reduction half-cell reaction must equal the number of electrons lost in the oxidation half-cell reaction.

The reactant which has undergone **reduction** caused the oxidation of another species and thus is termed the **oxidizing agent**.

The reactant which has undergone **oxidation** caused the reduction of another species and thus is termed the **reducing agent**.



Standard electrode potential (E° half-cell) is potential associated with a given half-cell reaction when all components are in standard states. The half-cell reaction is written as a reduction. The standard potential is also termed reduction potential. The chemical species closer to the top of the chart is more easily reduced, thus, a stronger oxidizing agent. The chemical species closer to the bottom of the chart is more easily oxidized, thus, a stronger reducing agent.



Predicting Cell Reactions

Is the reaction $\text{I}_2(\text{s}) + \text{Cu}(\text{s}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{I}^-(\text{aq})$ spontaneous?

(1) Determine the reduction potential values for the two half-cell reactions.



(2) Write oxidation and reduction half-cell reactions. Note the sign change for the oxidation half-cell reaction.



So this reaction is spontaneous, E°_{cell} has a positive value

Comparison of **Voltaic** versus **Electrolytic** Cells

Type of Cell	Voltaic Cell	Electrolytic Cell
Energy & Reaction Type	Energy is released from spontaneous redox reaction	Energy is absorbed to drive nonspontaneous redox reaction
System & Surroundings	System does work on load/ surroundings	Surroundings (power supply) does work on system (cell)
Diagram		
Oxidation = loss of e ⁻ Oxidation half-cell	occurs at anode $X \rightarrow X^+ + e^-$	occurs at anode $A^- \rightarrow A + e^-$
Reduction = gain of e ⁻ Reduction half-cell	occurs at cathode $e^- + Y^+ \rightarrow Y$	occurs at cathode $e^- + B^+ \rightarrow B$
Overall Reaction	$X + Y^+ \rightarrow X^+ + Y$ $\Delta G < 0$	$A^- + B^+ \rightarrow A + B$ $\Delta G > 0$
Electron flow	anode to cathode	anode to cathode
Charge on anode	negative	positive
Charge on cathode	positive	negative
Anions migrate	toward anode	toward anode
Cathode migrate	toward cathode	toward cathode
Requirements	need electrolyte (salt bridge, porous filter)	need electrolyte
Uses	fuel cells, batteries	plating, purifying active metals